## **Sugar-Sweetened Beverage Consumption**

**Source: Common Elements Questionnaire; The EARLY Trials** 

1. Over the <i>past 30 days</i> , how often did you drink soda or pop?	
	NEVER (Go to question 2)
	1 time per month or less
	2–3 times per month
	1–2 times per week
	3–4 times per week
	5–6 times per week
	1 time per day
	2–3 times per day
	4–5 times per day
	6 or more times per day
1a. How often were these sodas or pop diet or sugar-free?	
	Almost never or never
	About ¼ of the time
	About ½ of the time
	About ¾ of the time
	Almost always or always
2. Over the <i>past 30 days</i> , how often did you drink fruit drinks (such as cranberry cocktail, Hi-C, lemonade, or Kool-Aid, diet or regular)?	
	NEVER (GO TO QUESTION 3)
	1 time per month or less
	2–3 times per month
	1–2 times per week
	3–4 times per week
	5–6 times per week
	1 time per day
	2–3 times per day
	4–5 times per day
	6 or more times per day

2a. How often were your fruit drinks diet or sugar-free drinks?		
	Almost never or never	
	About ¼ of the time	
	About ½ of the time	
	About ¾ of the time	
	Almost always or always	
3. Over the <i>past 30 days</i> , how often did you drink sports drinks (such as Propel, PowerAde, or Gatorade)?		
	NEVER	
	1 time per month or less	
	2–3 times per month	
	1–2 times per week	
	3–4 times per week	
	5–6 times per week	
	1 time per day	
	2–3 times per day	
	4–5 times per day	
	6 or more times per day	
4. Over the <i>past 30 days</i> , how often did you drink energy drinks (such as Red Bull or Jolt)?		
	NEVER	
	1 time per month or less	
	2–3 times per month	
	1–2 times per week	
	3–4 times per week	
	5–6 times per week	
	1 time per day	
	2–3 times per day	
	4–5 times per day	
	6 or more times per day	